



Novel Approaches to the Analysis of  
Benzodiazepines and Morphine-Related Analytes  
in Forensic Samples

by  
Rebecca Webb

A thesis  
submitted for the degree of  
Doctor of Philosophy

Department of Chemistry, Materials and Forensic Science  
University of Technology, Sydney  
2005

Dedicated to the memory of Sir Joseph Swan (1828 – 1914) – pharmacist,  
physicist, inventor, chemist, and distant ancestor whose genetic influence has  
fuelled my passion for science.

### **Certificate of Authorship/Originality**

I certify that the work in this thesis has not previously been submitted for a degree nor has it been submitted as part of requirements for a degree.

I also certify that the thesis has been written by me. To the best of my knowledge, this thesis contains no material previously published or written by another person, except as fully acknowledged within the text. All information sources and literature used are indicated in the thesis. Any help that I have received in my research work and the preparation of the thesis itself has been acknowledged.

Rebecca Webb



---

## **Acknowledgements**

It's been a long, often difficult (but fulfilling) journey, made all the more easier thanks to a number of very special people who were always willing to offer me support and guidance when I most needed it. To my supervisors, Assoc Prof Michael Dawson and Dr Philip Doble – thank you for your ideas, encouragement and understanding. With your help, I have gained experience in so many different areas of research and the confidence to master any obstacle. I would also like to thank the staff from the Division of Analytical Laboratories (DAL), particularly Tania Prolov and Alan Hodda for facilitating my access to the necessary samples, and Jenny Easson for her assistance in the lab. Many thanks to all the staff and students in the Department of Chemistry, Materials and Forensic Science who helped me at various stages throughout my research; in particular, a very big thank you to Ronald Shimmon, Dr Kamali Kannangara, Dr Frank Cordaro, Mark Tahtouh and Tristan Rawling for their assistance with the synthetic aspects of my project. To all my fellow PhD students (past and present), especially Tam, Al, Katherine, Sonia, Matt, Jane, Daniel, Karen, Garry, Tristan, Mark – you've all been such great friends, and I know I would have gone crazy without you – thanks for all the good times! Finally, and most importantly, I have been lucky enough to have been blessed with wonderful supportive family and friends – too many to name individually, but you all know who you are. Dad, thanks for putting up with your “boomerang kid” for the past 18 months, and all the associated work! I'm forever grateful. Mum, you may live over 2000 km away, but you're still only a phone call or a short flight away. Thank you for always being there for me. To Jim – I would not be where I am today without you. You've helped me to realise what's important in life and what it takes to achieve it. Thank you for inspiring me every day of my life.



### List of Publications

Type of Publication	Number	Reference
Papers submitted to refereed journals	3	[1, 2, 3]
Poster presentations at International Meetings	4	[4, 5, 6, 7]

- [1] **Webb, R.**, Doble, P., Dawson, M., 2005, “A rapid capillary zone electrophoresis method for the analysis of benzodiazepines in spiked beverages”, submitted for publication in Journal of Forensic Sciences. (*Chapter 3*)
- [2] **Webb, R.**, Doble, P., Dawson, M., 2005, “Optimisation of HPLC Gradient Separations Using Artificial Neural Networks (ANNs): Application to Benzodiazepines in Post-Mortem Samples”, submitted for publication in Journal of Chromatography B. (*Chapter 4*)
- [3] **Webb, R.**, Dawson, M., Doble, P., Eason, J., Prolov, T., 2005, “Simultaneous Quantification of Morphine, Morphine-3-Glucuronide, Morphine-6-Glucuronide and Codeine in Post-Mortem Blood Samples”, submitted for publication in Journal of Forensic Sciences. (*Chapter 5*)
- [4] **Webb, R.**, Dawson, M., Kelly, T., Doble, P., 2002, “Analysis of morphine glucuronides”, ANZFSS 16<sup>th</sup> International Symposium on the Forensic Sciences, Canberra, May, 2002. (*Chapter 2*)
- [5] **Webb, R.**, Dawson, M., Kelly, T., Doble, P., 2002, “Analysis of morphine glucuronides”, Interact 2002, Sydney, July, 2002. (*Chapter 2*)

- [6] **Webb, R.**, Doble, P., Dawson, M., 2003, “Artificial neural networks for the optimisation of benzodiazepine analysis”, TIAFT 41<sup>st</sup> International Meeting, Melbourne, November, 2003. (*Chapter 4*)
- [7] **Webb, R.**, Doble, P., Dawson, M., 2003, “Artificial neural networks for the optimisation of benzodiazepine analysis”, ANZFSS 17<sup>th</sup> International Symposium on the Forensic Sciences, Wellington, March, 2004. (*Chapter 4*)

## List of Abbreviations

3MAM	3-monoacetylmorphine
6MAM	6-monoacetylmorphine
7-NH <sub>2</sub> -CLO	7-aminoclonazepam
7-NH <sub>2</sub> -FLU	7-aminoflunitrazepam
7-NH <sub>2</sub> -NIT	7-aminonitrazepam
Ac	acetyl
AC	acetylcodeine
ACN	acetonitrile
ADI	adinazolam
AIC	Australian Institute of Criminology
ALC	alcohol
ALP	alprazolam
ANN	artificial neural network
ANZFSS	Australia and New Zealand Forensic Science Society
APCI	atmospheric pressure chemical ionisation
B	blood
BBB	blood brain barrier
βCD	beta-cyclodextrin
BGE	background electrolyte
Bi	bile
BLOQ	below limit of quantification
BRO	brotizolam
BROM	bromazepam
bs	broad singlet
bt	broad triplet
C6G	codeine-6-glucuronide
CAR	carbamazepine
CD	cyclodextrin

CE	capillary electrophoresis
CEC	capillary electrochromatography
CEDIA	cloned enzyme donor immunoassay
CHL	chlordiazepoxide
C <sub>L</sub>	clearance
CLB	clobazam
CLO	clonazepam
CLOR	clorazepate
CLOT	clotiazepam
CLOX	cloxazolam
cm	centimetre
CMC	critical micelle concentration
CNS	central nervous system
COD	codeine
CSF	cerebrospinal fluid
CV	coefficient of variation
CZE	capillary zone electrophoresis
d	doublet
dt	double triplet
DAD	diode array detector
DAL	Division of Analytical Laboratories
DCM	dichloromethane
DEAE	<i>N</i> -diethylaminoethylene
DEL	delorazepam
DEM	demoxepam
DHC	dihydrocodeine
DHM	dihydromorphine
DIA	diazepam
DM $\beta$ CD	heptakis-(2,3-di- <i>O</i> -methyl)- $\beta$ -cyclodextrin
DS	dextran sulfate

E	erythrocytes
ECD	electrochemical detection
EIA	enzyme immunoassay
EKC	electrokinetic chromatography
ELISA	enzyme linked immunosorbent assay
EOF	electroosmotic flow
ESI	electrospray ionisation
ESP	electrospray
EST	estazolam
ETI	etizolam
EtOAc	ethyl acetate
F	fluorescence
FLU	flunitrazepam
FLUD	fludiazepam
FLUM	flumazenil
FLUR	flurazepam
FPIA	fluorescence polarisation immunoassay
GABA	gamma ( $\gamma$ )-aminobutyric acid
GC	gas chromatography
gluc	glucuronide
H	hair
HAL	halazepam
HALO	haloxazolam
HC	hydrocodone
HER	heroin
HM	hydromorphone
HPLC	high-performance liquid chromatography
hr or hrs	hour(s)
IPA	isopropyl alcohol
ISF	ideal separation function

$K_a$ (or $pK_a$ )	acid dissociation constant
KET	ketazolam
kg	kilogram
kV	kilovolts
L	litre
$L_{eff}$	effective length
LLE	liquid-liquid extraction
LOD	limit of detection
LOP	loprazolam
LOQ	limit of quantification
LOR	lorazepam
LORM	lormetazepam
LPME	liquid-phase microextraction
$L_{tot}$	total length
M	molar
m	multiplet
$m/z$	mass to charge ratio
M3G	morphine-3-glucuronide
M6G	morphine-6-glucuronide
$\mu_{app}$	apparent mobility
Mec	meconium
MED	medazepam
$\mu_{eff}$	effective mobility
MEKC	micellar electrokinetic chromatography
$\mu_{EOF}$	electroosmotic flow mobility
MeOH	methanol
MEX	mexazoram
mg	milligram
$\mu g$	microgram
MID	midazolam

min	minutes
mL	millilitre
MLP	multi-layer perceptron
mM	millimolar
mm	millimetre
µm	micrometre
mmol	millimole
MOR	morphine
MRM	multiple reaction monitoring
MS	mass spectrometry
MSA	methanesulfonic acid
mV	millivolts
ng	nanogram
-NH <sub>2</sub> -	amino
NIT	nitrazepam
nM	nanomolar
NMR	nuclear magnetic resonance spectroscopy
nor-CHL	norchlordiazepoxide
nor-CLB	norclobazam
nor-DIA	nordiazepam
nor-FLU	norflunitrazepam
nor-FLUD	norfludiazepam
NOS	noscaphine
OC	oxycodone
OD	overdose
-OH-	hydroxy
OXA	oxazepam
P	plasma
PAP	papaverine
PAR	paracetamol

PDA	photodiode array
PDDAC	Poly(diallyldimethylammonium chloride)
PRA	prazepam
PR <sub>s</sub>	product resolution
QSRR	quantitative structure-retention relationship
RBF	radial basis function
RIA	radioimmunoassay
$R_{\min}$	minimum resolution between consecutive peaks
RP	reversed phase
RRA	radioreceptor assay
S	serum
s	singlet
S/N	signal to noise ratio
SC	sodium cholate
SDC	sodium deoxycholate
SDS	sodium dodecyl sulphate
sec	seconds
SFE	supercritical fluid extraction
SIF	supervised injection facility
SIM	selected ion monitoring
SPE	solid-phase extraction
SPME	solid-phase microextraction
SRM	selected reaction monitoring
STA	systematic toxicological analysis
t	triplet
$t_{1/2}$	half-life
TCA	tricyclic antidepressant
TEA	triethylamine
TEAP	triethylammonium phosphate
TEM	temazepam



TET	tetrazepam
TFA	trifluoroacetic acid
THF	tetrahydrofuran
TIAFT	The International Association of Forensic Toxicologists
TLC	thin layer chromatography
TM $\beta$ CD	heptakis-(2,3,6-tri- <i>O</i> -methyl)- $\beta$ -cyclodextrin
TOF	tofisopam
TOF-SIMS	time of flight-secondary ion mass spectrometry
TRI	triazolam
TSP	thermospray
U	urine
UDP	uridine diphosphate
UDPGT	UDP-glucuronosyltransferases
UV	ultraviolet
V	volts
V <sub>D</sub>	volume of distribution
VH	vitreous humor

## Abstract

This research has focused on the development of analytical methods for the quantification of benzodiazepines and opiates in forensic samples. Both these classes of drugs are regularly encountered in forensic toxicology, especially in cases of fatal heroin overdose, and thus quantitative methods for their analysis are particularly relevant.

**Chapter 1** outlines the nature of the heroin problem in Australia and the different drug policies currently employed to help reduce the problems associated with heroin use and overdose. The involvement of metabolic factors and CNS depressants, such as benzodiazepines, in heroin overdose is discussed. An introduction to post-mortem drug analysis, as well as a comprehensive review of previously published methods for the analysis of heroin, its metabolites and benzodiazepines in biological samples, is presented.

**Chapter 2** describes procedures for the synthesis of commonly encountered morphine and benzodiazepine metabolites as an alternative to purchasing these metabolites commercially. Yields of 48, 25, 74 and 70% were obtained for M3G, M6G, 7-aminonitrazepam and 7-aminoclonazepam respectively.

**Chapter 3** documents the development of a rapid capillary zone electrophoresis (CZE) method for the analysis of nine benzodiazepines using dynamically prepared doubly-coated capillaries, consisting of a polycation of poly(diallyldimethylammonium chloride) (PDDAC) and a polyanion of dextran sulfate (DS). The addition of cyclodextrins to the background electrolyte (BGE) was also investigated as a means of improving analysis time. The validated method was successfully applied to the analysis of spiked beverages, with run times of less than 6.5 minutes.

**Chapter 4** describes the optimisation of a gradient HPLC separation of nine benzodiazepines using artificial neural networks (ANNs) in conjunction with experimental design. Various outputs and types of training data were investigated to yield the most appropriate ANN, which gave predictive errors of less than 5% for six of the nine analytes studied. A novel chromatographic function was also developed as a means of assessing the quality of chromatographic separations. The optimised method was validated for blood and successfully applied to authentic post-mortem samples. The limits of detection of the method ranged from 0.0057 to 0.023  $\mu\text{g/mL}$ , and recoveries were in the order of 58 – 92%.

**Chapter 5** details the development of a simple and rapid HPLC method for the simultaneous determination of morphine, M3G, M6G and codeine. Following SPE, the analytes were determined in post-mortem blood samples taken from heroin-related deaths. Cases involving the use of benzodiazepines in conjunction with heroin were found to have lower ratios of M6G/MOR and M3G/MOR, suggesting rapid death following heroin administration. M6G/M3G ratios were calculated to investigate the possible contribution of M6G towards heroin overdose with respect to its potential analgesic activity. M6G/M3G ratios were higher in cases involving the use of heroin only.

## Table of Contents

<b>Certificate of Authorship/Originality .....</b>	<b>iii</b>
<b>Acknowledgements .....</b>	<b>iv</b>
<b>List of Publications .....</b>	<b>v</b>
<b>List of Abbreviations .....</b>	<b>vii</b>
<b>Abstract.....</b>	<b>xiv</b>
<b>1 Introduction.....</b>	<b>1</b>
1.1 Summary .....	1
1.2 Origin of Heroin Production .....	2
1.3 Nature and Extent of the Heroin Problem.....	6
1.4 Drug Policy .....	8
1.5 Contributing Factors in Heroin Overdose.....	10
1.5.1 Hypersensitivity, Contamination and Batches.....	10
1.5.2 Systemic Dysfunction .....	11
1.5.3 Reduced Tolerance.....	11
1.5.4 Metabolic Variations.....	12
1.5.5 Drug Interactions .....	13
1.6 Post-mortem Drug Analysis.....	14
1.6.1 Specimen Selection.....	14
1.6.2 Extraction .....	15
1.6.3 Analysis.....	16
1.6.4 Post-mortem Considerations.....	16
1.7 Drugs of Interest .....	17
1.7.1 Heroin and Metabolites.....	17
1.7.1.1 Structural Features of Heroin and Related Compounds .....	17
1.7.1.2 Pharmacological Action and Effects.....	19
1.7.1.3 Metabolism and Pharmacokinetics .....	20
1.7.1.4 Analysis of Morphine and Metabolites in Biological Samples .....	23
1.7.1.5 Toxicological Data.....	34

1.7.2	Benzodiazepines .....	39
1.7.2.1	Structural Features of Benzodiazepines.....	39
1.7.2.2	Pharmacological Action and Effects.....	41
1.7.2.3	Metabolism and Pharmacokinetics .....	43
1.7.2.4	Analysis of Benzodiazepines in Biological Samples.....	47
1.7.2.5	Toxicological Data.....	68
1.8	Project Objectives .....	70
<b>2</b>	<b>Synthesis of Standard Drug Metabolites .....</b>	<b>73</b>
2.1	Summary .....	73
2.2	Introduction.....	73
2.2.1	Methods for the Synthesis of M3G, M6G, 7-Aminonitrazepam and 7-Aminoclonazepam .....	74
2.2.2	Objectives .....	77
2.3	Experimental .....	78
2.3.1	Chemicals.....	78
2.3.2	Instrumentation .....	79
2.3.3	Morphine-3-Glucuronide (M3G).....	80
2.3.4	Morphine-6-Glucuronide (M6G).....	80
2.3.4.1	Protection (I $\rightarrow$ V) .....	80
2.3.4.2	Condensation (V $\rightarrow$ VI).....	81
2.3.4.3	Deprotection (VI $\rightarrow$ VIII).....	82
2.3.5	7-Aminonitrazepam and 7-Aminoclonazepam .....	82
2.4	Conclusion .....	83
<b>3</b>	<b>Development of a Rapid Method for the Analysis of Benzodiazepines by Capillary Zone Electrophoresis (CZE) .....</b>	<b>86</b>
3.1	Summary .....	86
3.2	Introduction.....	86
3.2.1	Previous CE Methods for the Analysis of Benzodiazepines .....	90
3.2.2	Objectives .....	97
3.3	Experimental .....	97
3.3.1	Chemicals.....	97



3.3.2	Instrumentation .....	98
3.3.3	Calculations.....	99
3.3.4	Method Validation .....	100
3.4	Results and Discussion .....	100
3.4.1	Preliminary Screening of BGE Parameters.....	100
3.4.2	Coated Capillary Method Development .....	106
3.4.2.1	Stability of the EOF .....	115
3.4.3	Investigation of Cyclodextrins.....	116
3.4.4	Method Validation .....	123
3.4.5	Application of the Method .....	125
3.5	Conclusions.....	136
3.6	Future Work .....	136
<b>4</b>	<b>Optimisation of Chromatographic Separations Using Artificial Neural Networks (ANNs) and a Novel Chromatographic Function: Application to the Gradient HPLC Separation of Benzodiazepines.....</b>	<b>139</b>
4.1	Summary .....	139
4.2	Introduction.....	139
4.2.1	Artificial Neural Networks (ANNs).....	141
4.2.2	Previous Applications of Artificial Neural Networks to Gradient Elution Separations .....	145
4.2.3	Defining the Optimum Separation .....	146
4.2.4	Objectives .....	152
4.3	Experimental .....	152
4.3.1	Chemicals.....	152
4.3.2	Instrumentation .....	153
4.3.3	Calculations.....	153
4.3.4	Sample Preparation .....	154
4.3.5	Method Validation .....	155
4.3.6	Application of the Method .....	156
4.4	Results and Discussion .....	156
4.4.1	Preliminary Screening of Mobile Phase Parameters.....	156

4.4.2	ANN Modelling of the Experimental Space .....	164
4.4.2.1	Resolutions as the Output .....	168
4.4.2.2	Retention Times as the Output.....	174
4.4.3	Development of a Novel Function.....	187
4.4.4	Optimisation of the Separation Using ISF .....	198
4.4.5	Sample Preparation .....	204
4.4.6	Method Validation .....	207
4.4.7	Method Application .....	210
4.5	Conclusions.....	213
4.6	Future Work .....	214
<b>5</b>	<b>Simultaneous Quantification of Morphine, Morphine-3-Glucuronide, Morphine-6-Glucuronide and Codeine in Post-Mortem Blood Samples.....</b>	<b>216</b>
5.1	Summary .....	216
5.2	Introduction.....	216
5.2.1	Previous Investigations of M3G and M6G Concentrations and Ratios in Heroin Fatalities.....	219
5.2.2	Objectives .....	221
5.3	Experimental .....	222
5.3.1	Chemicals.....	222
5.3.2	Instrumentation .....	223
5.3.3	Calculations.....	223
5.3.4	HPLC Method Development .....	223
5.3.5	Sample Preparation .....	224
5.3.6	Method Validation .....	227
5.3.7	Method Application .....	228
5.4	Results and Discussion .....	228
5.4.1	HPLC Method Development .....	228
5.4.2	Sample Preparation .....	246
5.4.3	Method Validation .....	251
5.4.4	Method Application .....	254
5.5	Conclusions.....	265

5.6	Future Work .....	266
<b>6</b>	<b>Overall Conclusions .....</b>	<b>269</b>
<b>7</b>	<b>References .....</b>	<b>273</b>



## List of Figures

Figure 1-1 Synthesis of heroin from morphine.....	3
Figure 1-2 Geographical origin of heroin divided into a) Golden Crescent b) Golden Triangle c) Central and South America .....	5
Figure 1-3 Number of opioid overdose deaths in Australia among 15-44 yr olds, 1996-2001 .....	7
Figure 1-4 Some metabolic pathways of heroin .....	22
Figure 1-5 Metabolic pathways of some 1,4-benzodiazepines .....	45
Figure 2-1 Structure of acetobromo- $\alpha$ -D-glucuronic acid methyl ester (II).....	74
Figure 2-2 Synthesis of morphine-3-glucuronide (IV) .....	75
Figure 2-3 Synthesis of morphine-6-glucuronide (VIII).....	76
Figure 2-4 Synthesis of 7-aminoclonazepam (X) and 7-aminonitrazepam (XII) .....	77
Figure 3-1 Example electropherogram of a standard mixture of nine benzodiazepines on an uncoated capillary .....	102
Figure 3-2 Influence of phosphate buffer concentration on a) average effective mobilities and b) <b>average</b> resolution for each analyte at pH 2.5 on an uncoated capillary.....	104
Figure 3-3 Influence of pH on a) average effective mobilities and b) average resolution for each analyte in 100 mM phosphate buffer on an uncoated capillary.....	105
Figure 3-4 Influence of phosphate buffer concentration on a) average effective mobilities and b) average resolution for each analyte at pH 2.5 on a coated capillary.....	108
Figure 3-5 Influence of pH on a) average effective mobilities and b) average resolution for each analyte in 100 mM phosphate buffer on a coated capillary .....	109
Figure 3-6 Example electropherogram of a standard mixture of nine benzodiazepines on a coated capillary .....	111
Figure 3-7 Influence of a) phosphate buffer concentration and b) pH on average product resolution for coated and uncoated capillaries.....	112

Figure 3-8 Influence of a) phosphate buffer concentration and b) pH on run time for coated and uncoated capillaries .....	113
Figure 3-9 Effect of DS in running buffer on the coefficient of variation (%) of the EOF .....	116
Figure 3-10 Influence of cyclodextrin type and concentration on product resolution (PR <sub>s</sub> ) for a coated capillary .....	118
Figure 3-11 Influence of cyclodextrin type and concentration on product resolution (PR <sub>s</sub> ) for an uncoated capillary .....	119
Figure 3-12 Maximum product resolution (PR <sub>s</sub> ) attained for each type of cyclodextrin on coated and uncoated capillaries .....	121
Figure 3-13 Run times for each type of cyclodextrin on coated and uncoated capillaries .....	121
Figure 3-14 Example electropherograms of a) spiked Bacardi and b) blank Bacardi .....	128
Figure 3-15 Example electropherograms of a) spiked bourbon and b) blank bourbon .....	129
Figure 3-16 Example electropherograms of a) spiked Coca-cola and b) blank Coca-cola.....	130
Figure 3-17 Example electropherograms of a) spiked wine and b) blank wine .....	131
Figure 3-18 Example electropherograms of a) spiked beer and b) blank beer .....	132
Figure 3-19 Example electropherograms of a) spiked orange juice and b) blank orange juice .....	133
Figure 4-1 Example of ANN architecture (MLP 3:3-4-2:2).....	142
Figure 4-2 Influence of pH on a) average retention times and b) average resolution for each analyte .....	158
Figure 4-3 Influence of formate buffer concentration on a) average retention times and b) average resolutions for each analyte .....	159
Figure 4-4 Influence of methanol composition on a) average retention times and b) average resolutions for each analyte .....	161
Figure 4-5 Influence of acetonitrile composition on a) average retention times and b) average resolutions for each analyte .....	162

Figure 4-6 Effect on elution order of a) ACN and b) MeOH .....	163
Figure 4-7 Example of percentage composition of each mobile phase component with varying time (15% MeOH, 2%/min ACN, 5% initial ACN).....	165
Figure 4-8 Schematic of experimental space .....	167
Figure 4-9 Overall correlation between experimental and predicted peak pair resolutions for a 3:3-12-8:8 MLP ANN trained on replicate data .....	169
Figure 4-10 Overall correlation between experimental and predicted peak pair resolutions for a 3:3-12-8:8 MLP ANN trained on average data.....	170
Figure 4-11 Overall correlation between experimental and predicted retention times using a 3:3-20-9:9 MLP ANN trained on replicate data.....	175
Figure 4-12 Overall correlation between experimental and predicted retention times using a 3:3-12-9:9 MLP ANN trained on average data .....	176
Figure 4-13 Overall correlation between experimental and predicted peaks widths for a 3:3-12-9:9 MLP ANN .....	182
Figure 4-14 Influence of initial %ACN and ACN gradient product resolution (at 10% MeOH).....	185
Figure 4-15 Influence of %MeOH and initial %ACN product resolution (at ACN gradient 1) .....	185
Figure 4-16 Influence of ACN gradient and %MeOH on product resolution (at 5% initial ACN) .....	186
Figure 4-17 Example chromatogram of the separation optimised using product resolution.....	186
Figure 4-18 Range and domain for the resolution term of the ISF .....	195
Figure 4-19 Range and domain for the run time term of the ISF .....	196
Figure 4-20 Range and domain for the void term of the ISF.....	197
Figure 4-21 ISF response surfaces for varying a) %MeOH and ACN gradient, b) initial %ACN and ACN gradient and c) initial %ACN and %MeOH.....	200
Figure 4-22 Example chromatogram of the optimised separation of nine benzodiazepines .....	202
Figure 4-23 Comparison of average recovery obtained for ethyl acetate, diethyl ether/ethyl acetate and diethyl ether LLEs.....	205



Figure 4-24 Example chromatograms of a) blank and b) spiked whole blood extracted with diethyl ether/ethyl acetate (1:1).....	207
Figure 4-25 Example chromatograms for post-mortem blood samples testing positive for benzodiazepines a) case 2401190 b) case 2402494 .....	211
Figure 5-1 Effect of various excitation and emission wavelengths on the signal/noise ratio of M6G .....	229
Figure 5-2 Comparison of signal/noise ratios obtained for each analyte using UV and fluorescence detection.....	229
Figure 5-3 Effect of organic modifier type on retention times of each analyte .....	230
Figure 5-4 Effect of varying formate buffer pH on a) average retention times and b) average resolution for each analyte on the Zorbax SB-C <sub>18</sub> stationary phase...	233
Figure 5-5 Effect of varying acetate buffer pH on a) average retention times and b) average resolution for each analyte on the Zorbax SB-C <sub>18</sub> stationary phase...	234
Figure 5-6 Effect of varying buffer concentration on a) average retention times and b) average resolution of each analyte on the Zorbax SB-C <sub>18</sub> stationary phase	236
Figure 5-7 Effect of varying temperature on a) average retention times and b) average resolution of each analyte on the Zorbax SB-C <sub>18</sub> stationary phase ....	237
Figure 5-8 Effect of varying formate buffer pH on a) average retention times and b) average resolution for each analyte on the Atlantis dC <sub>18</sub> stationary phase .....	240
Figure 5-9 Effect of varying acetate buffer pH on a) average retention times and b) average resolution for each analyte on the Atlantis dC <sub>18</sub> stationary phase .....	241
Figure 5-10 Effect of varying concentration on a) average retention times and b) average resolution for each analyte on the Atlantis dC <sub>18</sub> stationary phase .....	242
Figure 5-11 Effect of varying temperature on a) average retention times and b) average resolution of each analyte on the Atlantis dC <sub>18</sub> stationary phase .....	243
Figure 5-12 Example chromatogram of the separation of M3G, M6G, morphine, codeine and 6MAM .....	245
Figure 5-13 Comparison of recoveries of each analyte using precipitation and LLE techniques .....	248
Figure 5-14 Comparison of recoveries of each analyte using SPE techniques.....	249

Figure 5-15 Example chromatogram of spiked whole blood extracted using Bond Elut C <sub>18</sub> SPE.....	251
Figure 5-16 Relationship between the free morphine concentration and the ratio of a) M6G/MOR b) M3G/MOR in heroin and heroin/benzodiazepine fatalities.....	260
Figure 5-17 Relationship between M6G/MOR and M6G/M3G in heroin and heroin/benzodiazepine fatalities.....	262
Figure 5-18 Example chromatogram of a post-mortem blood sample from a heroin-related death (Case 2402603).....	263

## List of Tables

Table 1-1 Structures of some compounds chemically related to heroin.....	18
Table 1-2 SPE cartridges used for the extraction of heroin metabolites from biological samples.....	24
Table 1-3 Summary of published HPLC methods for heroin metabolites in biological samples.....	26
Table 1-4 Free and total morphine concentrations and ratios in fatal heroin overdose victims and survivors .....	36
Table 1-5 Concentration ranges and molar ratios for MOR, M3G and M6G in heroin fatalities.....	37
Table 1-6 Concentration ranges and molar ratios for MOR, M3G and M6G in morphine therapy patients.....	38
Table 1-7 Structures of selected 1,4-benzodiazepines .....	40
Table 1-8 Structures of selected imidazo- and triazolo-benzodiazepines.....	41
Table 1-9 Pharmacokinetic parameters of some benzodiazepines.....	47
Table 1-10 SPE cartridges used for the extraction of benzodiazepines from biological samples.....	48
Table 1-11 Summary of published HPLC methods for benzodiazepines in biological samples.....	51
Table 1-12 Summary of published CE methods for benzodiazepines in biological samples.....	65
Table 1-13 Therapeutic and toxic ranges of benzodiazepines in serum .....	69
Table 1-14 Fatal concentration ranges of selected benzodiazepines in blood .....	69
Table 3-1 Summary of published CE methods for the analysis of benzodiazepines.	92
Table 3-2 Average resolutions of consecutive peaks with varying phosphate buffer concentration on an uncoated capillary.....	104
Table 3-3 Average resolutions of consecutive peaks with varying pH on an uncoated capillary.....	105
Table 3-4 Average resolutions of consecutive peaks with varying phosphate buffer concentration on a coated capillary.....	108

Table 3-5 Average resolutions of consecutive peaks with varying pH on a coated capillary.....	109
Table 3-6 Average resolutions for consecutive peaks with varying type and concentration of cyclodextrin on a coated capillary .....	118
Table 3-7 Average resolutions for consecutive peaks with varying type and concentration of cyclodextrin on an uncoated capillary .....	119
Table 3-8 Calibration and LOD data.....	124
Table 3-9 Accuracy and precision data.....	124
Table 3-10 Recovery of each drug from spiked beverages.....	134
Table 4-1 Optimisation functions for assessing chromatographic separations.....	149
Table 4-2 Evaluated LLE techniques for benzodiazepine analytes .....	154
Table 4-3 Average resolutions of consecutive peaks with varying pH.....	158
Table 4-4 Average resolutions of consecutive peaks with varying formate buffer concentration.....	159
Table 4-5 Average resolutions of consecutive peaks with varying methanol composition.....	161
Table 4-6 Average resolutions of consecutive peaks with varying acetonitrile composition.....	162
Table 4-7 Values for each experiment in three factor design .....	167
Table 4-8 Correlation equations and correlation coefficients between experimental and predicted resolutions for each peak pair trained on replicate data .....	169
Table 4-9 Correlation equations and correlation coefficients between experimental and predicted resolutions for each peak pair trained on average data .....	170
Table 4-10 Experimental and predicted resolution values for each ANN with corresponding relative errors .....	173
Table 4-11 Correlation equations and correlation coefficients between experimental and predicted retention times for each analyte trained on replicate data.....	175
Table 4-12 Correlation equations and correlation coefficients between experimental and predicted retention times for each analyte trained on average data .....	176
Table 4-13 Experimental and predicted retention times for each ANN with corresponding relative errors .....	179



Table 4-14 Correlation equations and correlation coefficients between experimental and predicted peak widths for each analyte .....	182
Table 4-15 Experimental and predicted peak widths with corresponding relative errors .....	183
Table 4-16 Resolutions between consecutive peaks at the optimum predicted by PR <sub>s</sub> .....	186
Table 4-17 Experiments selected for the comparison of optimisation functions.....	188
Table 4-18 Comparison of selected optimisation functions for evaluating separations .....	191
Table 4-19 Assessment of separations according to the ISF .....	197
Table 4-20 Predicted and experimental retention times and peak widths with corresponding errors for the optimum separation conditions .....	202
Table 4-21 Predicted and experimental resolution values for the optimum separation conditions .....	202
Table 4-22 Calibration and LOD data for benzodiazepine analytes in blood.....	208
Table 4-23 Intra- and inter-assay precision and accuracy data for benzodiazepine analytes in blood .....	209
Table 4-24 Blood concentrations of benzodiazepines in post-mortem blood samples .....	210
Table 5-1 Evaluated LLE techniques for the extraction of morphine and related analytes .....	225
Table 5-2 Evaluated SPE methods for the extraction of morphine and related analytes .....	226
Table 5-3 Average resolution of each analyte pair with varying pH of formate buffer on the Zorbax SB-C <sub>18</sub> stationary phase.....	233
Table 5-4 Average resolution of each analyte pair with varying pH of acetate buffer on the Zorbax SB-C <sub>18</sub> stationary phase.....	234
Table 5-5 Average resolution of each analyte pair with varying buffer concentration on the Zorbax SB-C <sub>18</sub> stationary phase.....	236
Table 5-6 Average resolution of each analyte pair with varying temperature on the Zorbax SB-C <sub>18</sub> stationary phase.....	237



Table 5-7 Average resolution of each analyte pair with varying formate buffer pH on the Atlantis dC <sub>18</sub> stationary phase.....	240
Table 5-8 Average resolution of each analyte pair with varying acetate buffer pH on the Atlantis dC <sub>18</sub> stationary phase.....	241
Table 5-9 Average resolution of each analyte pair with varying concentration on the Atlantis dC <sub>18</sub> stationary phase.....	242
Table 5-10 Average resolution of each analyte pair with varying temperature on the Atlantis dC <sub>18</sub> stationary phase.....	243
Table 5-11 Recoveries of each analyte using precipitation and LLE techniques ....	248
Table 5-12 Recoveries of each analyte using SPE techniques.....	249
Table 5-13 Calibration and LOD data for morphine and related analytes in blood	253
Table 5-14 Intra- and inter-assay precision and accuracy data for morphine and related analytes in blood .....	253
Table 5-15 Molar ratios and blood concentrations of morphine, M3G, M6G and codeine in heroin fatalities .....	255
Table 5-16 Molar ratios and blood concentrations of morphine, M3G, M6G, codeine and benzodiazepines in heroin/benzodiazepine fatalities .....	256

## List of Equations

Equation 3-1 Calculation of apparent mobility ( $\mu_{app}$ ).....	99
Equation 3-2 Calculation of effective mobility ( $\mu_{eff}$ ).....	99
Equation 3-3 Calculation of resolution ( $R_s$ ).....	99
Equation 3-4 Calculation of product resolution ( $PR_s$ ) .....	100
Equation 4-1 Calculation of resolution ( $R_s$ ).....	153
Equation 4-2 Calculation of product resolution ( $PR_s$ ) .....	153
Equation 4-3 Calculation of relative error ( $\%e_i$ ) for resolution .....	154
Equation 4-4 Calculation of relative error ( $\%e_i$ ) for retention times .....	154
Equation 4-5 Calculation of normalised product resolution .....	189
Equation 4-6 Calculation of CRS .....	189
Equation 4-7 Calculation of CRF .....	189
Equation 4-8 Calculation of CEF.....	189
Equation 4-9 Calculation of Q .....	190
Equation 4-10 The Ideal Separation Function (ISF).....	194
Equation 5-1 Calculation of resolution ( $R_s$ ).....	223

# **Chapter One**

## **Introduction**